

Introduction to climate modelling

Steven J. Phipps


Climate Change Research Centre
ARC Centre of Excellence for Climate System Science
University of New South Wales



`s.phipps@unsw.edu.au`


Overview

1. Why do we need climate models?
2. What is a climate model?
3. How does a climate model work?
4. How do you build a climate model?
5. How do you use a climate model?
6. Examples of climate modelling



1. Why do we need climate models?







Northern Hemisphere



Why do we need climate models?


- There is only one Earth, and we can't (shouldn't) perform experiments on that
- We can't travel in time
- We want to predict possible future climate states
- We want to understand past climatic changes
- We want to explore properties of the climate system
- We want to answer *questions* - these can range from *scientific* questions to *policy* questions

2. What is a climate model?

- A *virtual* Earth
- A computer program (usually very long and complex)
- Solves the fundamental physical equations that describe the evolution of the climate system
- Divides the Earth into discrete components
- Different types of models: simple vs. complex, low-resolution vs. high-resolution, regional vs. global
- A model is a *tool* - the type that you use depends upon the question that you want to answer
- *No* model is a perfect representation of the real world

A virtual Earth

Modeling the Climate System



A computer program

```
c... Calculate density
      p2 = p*p
      p3 = p*p2
      do i = 1, imt
        s15(i) = s(i)*sqrt(s(i))
        s2(i) = s(i)*s(i)
        t2(i) = t(i)*t(i)
        t3(i) = t(i)*t2(i)
        t4(i) = t(i)*t3(i)
        rho(i) = (a0 + a1*t(i) + a2*t2(i) + a3*t3(i) + a4*s(i) +
&                a5*s(i)*t(i) + a6*s2(i) + a7*p + a8*p*t2(i) +
&                a9*p*s(i) + a10*p2 + a11*p2*t2(i)) /
&                (b0 + b1*t(i) + b2*t2(i) + b3*t3(i) + b4*t4(i) +
&                b5*s(i) + b6*s(i)*t(i) + b7*s(i)*t3(i) + b8*s15(i) +
&                b9*s15(i)*t2(i) + b10*p + b11*p2*t3(i) + b12*p3*t(i))
      end do
```

Solves fundamental physical equations


$$\rho(S, \theta, p) = \frac{P_1(S, \theta, p)}{P_2(S, \theta, p)} \quad (1)$$


where


$$\begin{aligned} P_1(S, \theta, p) = & a_0 + a_1\theta + a_2\theta^2 + a_3\theta^3 + a_4S \\ & + a_5S\theta + a_6S^2 + a_7p + a_8p\theta^2 \\ & + a_9pS + a_{10}p^2 + a_{11}p^2\theta^2 \end{aligned} \quad (2)$$

$$\begin{aligned} P_2(S, \theta, p) = & b_0 + b_1\theta + b_2\theta^2 + b_3\theta^3 + b_4\theta^4 \\ & + b_5S + b_6S\theta + b_7S\theta^3 + b_8S^{\frac{3}{2}} \\ & + b_9S^{\frac{3}{2}}\theta^2 + b_{10}p + b_{11}p^2\theta^3 + b_{12}p^3\theta \end{aligned} \quad (3)$$

Divides the Earth into discrete components










Figure 53. A schematic representation of the essential components of a fully coupled general circulation model, based on the conservation of mass, energy and momentum in the atmosphere and ocean, and the physical processes involved in the coupling between them.

Different types of models




Fig. 1. Pictorial definition of EMICs. Adapted from Claussen (2000)





Fig. 2. The climate modeling pyramid. Adapted from Henderson-Sellers and McGuffie (1987)


The Development of Climate models, Past, Present and Future



Box 3, Figure 1: The development of climate models over the last 25 years showing how the different components are first developed separately and later coupled into comprehensive climate models.


Limitations on the accuracy of climate models

- Lack of understanding of physical processes: if we don't understand a process, we can't describe it within a model
- Computational limitations:
 - It's impossible to include all physical processes in a single model, so some processes are always missing
 - Limited spatial resolution
- Essential to comprehensively evaluate a model before trusting the output



Models work!


- The figures on the right show annual rainfall
- Which is observed and which is modelled?



3. How does a climate model work?

- A climate model accepts input data:
 - Initial conditions
 - Boundary conditions
- Iterates forwards in time, typically using an interval of 15 minutes to 1 hour
- Starting from the initial state of the climate system, applies physical laws to calculate the state of the climate system at the next time interval
- Repeats this process for as long as necessary
- Generates output data (lots!)

Climate Modelling



4. How do you build a climate model?

- Identify the processes to be modelled
- Identify the quantities to be modelled
- Identify the relationships between these quantities
- Express these relationships mathematically
- Write computer code to solve these equations

How do you build a climate model?

- Traditional approach:
 - Develop a computer program from scratch
- Modern approach:
 - Take existing components and combine them
- Test and debug
- Determine the optimal parameter settings (“tuning”)
- Evaluate, evaluate, evaluate...


How do you build a climate model?

- *Very* specialised and time-consuming process
- The end result is a very large and complex computer program
- A typical state-of-the-art climate model:
 - represents *hundreds* of person-years of work
 - consists of hundreds of thousands, or even millions, of lines of computer code
 - is very computationally expensive
 - generates enormous amounts of data

Case study: ACCESS

- Australian Community Climate and Earth System Simulator
- Atmosphere: Unified Model (UK)
- Ocean: MOM4 (USA)
- Sea ice: CICE (USA)
- Land surface: CABLE (Australia)
- Coupler: OASIS (France)
- Around one million lines of source code
- Can simulate around 2-3 years per day
- Generates up to 50 GB of data for each year

ACCESS ESM





5. How do you use a climate model?

- Select the question that you want to answer
- Select an appropriate model
- Configure the model accordingly:
 - Initial conditions
 - Boundary conditions
- Find a big enough computer, and somewhere to store the data...


National Facility, Canberra








National Facility, Canberra




6. Examples of climate modelling









(a) Mediaeval Climate Anom (O+G)




(d) Little Ice Age (O+G)




(g) MCA minus LIA (O+G)




(b) Mediaeval Climate Anom (O+G+S)




(e) Little Ice Age (O+G+S)




(h) MCA minus LIA (O+G+S)



(c) Mediaeval Climate Anom (O+G+S+V)



(f) Little Ice Age (O+G+S+V)



(i) MCA minus LIA (O+G+S+V)